

A new universality class describing the insulating regime of disordered wires with strong spin-orbit scattering

M. Caselle

*Dipartimento di Fisica Teorica dell'Università di Torino and I.N.F.N., Sezione di Torino
via P.Giuria 1, I-10125 Turin, Italy*

We discuss the distribution of transmission eigenvalues in the strongly localized regime in the presence of both a magnetic field and spin-orbit scattering. We show that, under suitable conditions, this distribution can be described by a new universality class labelled not only by the index β but also by a new index η . This result is obtained by mapping the problem into that of a suitable Calogero-Sutherland model.

PACS numbers: 72.10.Bg, 05.60.+w, 72.15.Rn, 72.20.My

Magnetoconductance (MC) in the strongly localized regime of mesoscopic systems has been the subject of an increasing interest during last years [1] - [5]. While the same phenomenon in the weakly localized regime is by now rather well understood, and a good agreement has been reached among different theoretical descriptions and between their predictions and the experimental results, the situation is less clear in the strongly localized regime. In particular it is not clear to which extent the Random Matrix Theory (RMT) of quantum transport [6,7], and the corresponding classification in terms of universality classes, originally developed to describe the weakly localized regime, can also be extended to the insulating regime. Indeed, it seems that two different insulating regimes exist. The first one (strongly insulating regime) in which the localization length ξ (to be defined below) is much smaller than the hopping length t_h , seems to be well described by a directed-path approach (see ref. [1,5] and references therein). The second one (weakly insulating regime), close to the Anderson transition, in which $t_h/\xi \sim 1$ and backscattering paths are important, seems to be well described within a RMT framework. This would be an important result due to the simplicity and generality of the RMT approach: the input required is only the symmetry of the Hamiltonian and the resulting classification is completely encoded by a single index β . The above scenario is however complicated by the presence of some conflicting theoretical analyses. The disagreement between the various approaches becomes evident when looking at the MC behaviour in presence of strong spin orbit scattering (SOS). In fact in the insulating regime the localization length is related to the conductance by: $g \equiv \exp(-2L/\xi)$, (L =length of the sample). In the standard RMT approach ξ is proportional to β and, as a consequence, the magnetoconductance is directly related to the change of β as a function of the magnetic field. The RMT approach based on the maximal entropy ansatz predicts that the effect of a magnetic field on a sample with SOS is to change β from $\beta = 4$ ("symplectic ensemble": spin-rotation symmetry broken but time reversal symmetry conserved) to $\beta = 2$ ("uni-

tary ensemble": time reversal symmetry broken, status of the spin-rotation symmetry irrelevant). Defining as $\xi(H)$ the localization length in presence of a magnetic field, from the above analysis it follows that $\xi(H)/\xi = 1/2$, and we expect a strong *negative* MC [3]. On the contrary an exact calculation within the non-linear σ model [8] shows that if the spin-rotation symmetry is broken, β remains equal to 4 even in presence of a breaking of the time reversal symmetry (thus implying a zero MC) and that in this case the ensemble behaves as unitary ($\beta = 2$) only with respect to the level statistics. In this letter we shall try to clarify this issue. Let us state our main results:

a] The main point we want to stress is that, while in the weakly localized regime all the relevant physical properties are completely determined by the level statistics (namely the value of β), when extending the RMT approach to the insulating regime, due to the fact that in this case *the conductance is dominated by the lowest eigenvalue* all the details of the chosen RMT model (number of degrees of freedom, possibly the presence of new critical indices) become important.

b] In particular this gives a way to reconcile the above described controversy. We shall show below that also within the maximal entropy approach a zero MC should be expected in the insulating regime.

c] It is however very interesting to notice that in some recent experiments [4] on samples characterized by strong SOS a clear negative MC has been observed in the weakly insulating regime, in contradiction with the above results. We shall show below that under suitable hypothesis, it does exist a new universality class in the RMT description of quantum transport which indeed admits negative MC in the insulating regime. This new class has $\beta = 4$, as in [8], but a new index η exists that allows to distinguish this case from the ordinary symplectic ensemble. As a consequence of this new η index a negative MC is expected even if $\beta = 4$.

To show these results let us first briefly remind few known features of the RMT description of quantum transport. The central object in this approach is the

$2N \times 2N$ Transfer Matrix M (where N is the number of scattering channels at the Fermi level) which relates the incoming and outgoing fluxes in a two-probe geometry (namely a finite disordered section of length L and transverse width L_t , to which current is supplied by two semi-infinite ordered leads):

$$M \begin{pmatrix} I \\ O \end{pmatrix} = \begin{pmatrix} O' \\ I' \end{pmatrix}, \quad (1)$$

where the I, O, I', O' vectors describe the incoming and outgoing wave amplitudes on the left and right respectively. To construct the RMT one must first impose the physical requirements which define the ensemble of the matrices: for instance flux conservation, or time conjugation, if time reversal symmetry is conserved. This defines the group G to which the random matrix belongs. Then one assumes that the physics of the problem only depends on a subset of the degrees of freedom of the matrix. It turns out that a good parametrization in our case is given by the N real, positive eigenvalues λ_i of the matrix $Q = \frac{1}{4}[M^\dagger M + (M^\dagger M)^{-1} - 2]$. These parameters have an important physical interpretation since they are related to the transmission eigenvalues T_i by $\lambda_i \equiv (1 - T_i)/T_i$ in terms of which all the interesting physical properties can be evaluated. For instance the conductance g is given by $g = g_0 \sum_n T_n$ (with $g_0 = 2e^2/h$). This parametrization allows to identify a subgroup H of invariance in our ensemble. The matrix ensemble is thus related to a suitable G/H coset. It turns out that in all the relevant cases G/H is actually a symmetric space. The last step is the identification of a dynamical principle to describe the “time” evolution of the eigenvalues. This will allow to construct explicitly the probability distribution $P(\{\lambda_i\})$ for the eigenvalues λ_i . In the case of RMT for quantum transport the “time” $s = L/l$ is actually the length of the conductor L measured in unities of the mean free path l ; the dynamical principle is the so called “local maximum entropy principle” which simply implies the free motion of the eigenvalues in the symmetric space G/H and, as a consequence, the evolution operator is the radial part of the Laplace-Beltrami operator on G/H . The corresponding (coupled) differential equation for the probability distribution $P(\lambda)$ of the eigenvalues is known as the Dorokhov-Mello-Pereyra-Kumar (DMPK) equation [9]

$$\frac{\partial P}{\partial s} = \frac{2}{\gamma} \sum_{i=1}^N \frac{\partial}{\partial \lambda_i} \lambda_i (1 + \lambda_i) J \frac{\partial}{\partial \lambda_i} J^{-1} P \equiv D P, \quad (2)$$

where $\gamma = \beta N + 2 - \beta$ and $\beta \in \{1, 2, 4\}$ is the symmetry index discussed above. $J(\{\lambda_n\})$ denotes the Jacobian from the matrix to the eigenvalue space:

$$J(\{\lambda_n\}) = \prod_{i < j} |\lambda_j - \lambda_i|^\beta. \quad (3)$$

We have seen that the RMT is completely identified by its G/H symmetric space. All the irreducible symmetric

spaces of classical type can be classified with essentially the same techniques used for the Lie algebras. They fall into 11 classes labelled by the type of root system and by the multiplicities of the various roots [10]. The DMPK operator D is completely fixed once the information on the type of root system and the root multiplicities are given. The Dyson ensembles, which first appeared in the RMT of nuclear energy levels, (and are also relevant in the RMT of quantum transport based on the scattering matrix [11]) are related to the symmetric spaces labelled by the Dynkin diagrams of type A_N . There are exactly three classes of A_N spaces, which are distinguished by the index β which labels the multiplicity of the roots. β can take the values 1, 2 or 4 and exactly coincides with the symmetry index of the corresponding Random Matrix ensemble. Instead, the ensembles or random transfer matrices in which we are interested coincide with the symmetric spaces labelled by Dynkin diagrams of type C_N (see Tab.1). *It is exactly the difference between A_N and C_N type of symmetric spaces the origin of the differences between RMT based on transfer matrices and the Wigner-Dyson ones.*

G	H	d	β	η
$Sp(2N, \mathbf{R})$	$U(N)$	$N(N+1)$	1	1
$SU(N, N)$	$SU(N) \otimes SU(N) \otimes U(1)$	$2N^2$	2	1
$SO^*(4N)$	$U(2N)$	$N(N-1)$	4	1
$USp(2N, 2N)$	$USp(2N) \otimes USp(2N)$	$4N^2$	4	3
$Sp(2N, \mathbf{C})$	$USp(2N)$	$N(2N+1)$	2	2

TABLE I. Irreducible symmetric spaces of type C_N . In the first two columns the group G and the maximal subgroup H which define the symmetric space. In the third column the dimensions d of the space. All these spaces are labelled by the Dynkin diagram C_N . In the fourth column the multiplicity β of the ordinary roots of C_N . In the last column the multiplicity η of the long root.

The most relevant feature of the C_N symmetric spaces with respect of the A_N ones is that they are described by two critical indices instead of one. This is due to the fact that the C_N Dynkin diagrams have two types of roots, each with its particular multiplicity. We shall call this second index η in the following. This peculiar feature of these ensembles was up to now ignored because in the weakly localized regime it is only the value of β which matters, and also because the three cases which have been studied up to now (the first three lines of tab.1) have the same value of the second index $\eta = 1$, thus leading to the same DMPK equation (with no explicit η dependence). However we shall show now that in the insulating regime, if *both* the spin-rotation and the time reversal symmetry are broken, also the fourth ensemble of those listed in tab.1 could become relevant, and in this case the value of η changes.

As a preliminary step to prove this assertion, let us remind the group properties of the symplectic ensemble of transfer matrices (spin-rotation broken, time reversal conserved). In this case each one of the I, I', O, O' vectors is a collection of N spinors (one for each channel); each spinor contains the two spin degrees of freedom. Hence M is in this case a $4N \times 4N$ complex matrix.

Flux conservation implies

$$|I|^2 - |O|^2 = |O'|^2 - |I'|^2 \quad , \quad (5)$$

which can be written as:

$$M^\dagger \Sigma_z M = \Sigma_z \quad (6)$$

where

$$\Sigma_z = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix} \quad (7)$$

where $\mathbf{1}$ and $\mathbf{0}$ denote the $2N \times 2N$ unit and zero matrix. Eq.(6) implies that the matrix M belongs to the unitary group $U(2N, 2N)$ group (since it preserves the metric Σ_z). Time reversal must then be imposed in two steps. At the level of the single spinor we must require invariance with respect to the time reversal operator: $\theta = i\sigma_y C$ where C is the complex conjugation operator and σ_y is the Pauli matrix:

$$\theta = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} C \quad (8)$$

In addition, at the level of the whole matrix M one must impose the following constraint

$$M^* = \Sigma_x M \Sigma_x \quad (9)$$

with

$$\Sigma_x = \begin{pmatrix} \mathbf{0} & \mathbf{1} \\ \mathbf{1} & \mathbf{0} \end{pmatrix} \quad (10)$$

which, together with eq.(6) implies that $M \in SO^*(4N)$. Let us now switch on the magnetic field. We have already seen that if both the constraints (8) and (9) are eliminated we have $M \in U(2N, 2N)$, namely that M belongs in this case to the unitary ensemble $\beta = 2$ listed in tab.1, *but with* $N \rightarrow 2N$. This gives the standard RMT prediction: β changes from 4 to 2 in relation to the level statistics, as a consequence of the application of a magnetic field. However for all those observables which are not only related to the level statistics, but depend more generally on γ , the two changes $\beta : 4 \rightarrow 2$, $N \rightarrow 2N$ compensate and the model effectively behaves as if $\beta = 4$ as in the σ model result. This happens in particular for the localization length ξ in the insulating regime in which we are interested. In fact we shall show below that ξ is proportional to γ . As mentioned above this seems to exclude a negative MC in the RMT approach, in clear

contradiction to the experimental results. To open the possibility to find a negative MC with RMT, we must make a further hypothesis. Since the two constraint (8) and (9) are not on the same footing, the first being a local and the second a global property of the electrons, they need not to be simultaneously fulfilled. In particular it could exist a range of parameters (say, for low magnetic fields) in which only the second one is eliminated but the first one survives. Physically this would mean that locally the Kramers degeneracy is conserved, but globally time-reversal is broken, with all the known consequences on the back-scattering paths. Let us see which are the consequences of this hypothesis. It is easy to see that eq.(8) alone implies the conservation of the antisymmetric metric and that in this case the joint application of (6) and (8) leads to $M \in USp(2N, 2N)$. This choice of constraints defines the new universality class anticipated in the title. Since $USp(2N, 2N) = U(N, N, \mathbf{Q})$ we see that in this case M has a very natural description in terms of quaternions, being the most general unitary quaternion matrix which preserves the metric Σ_z . Any matrix M of this type can be parametrized as:

$$M = \begin{pmatrix} u^{(1)} & \mathbf{0} \\ \mathbf{0} & u^{(2)} \end{pmatrix} \begin{pmatrix} \sqrt{1+\mathbf{\Lambda}} & \sqrt{\mathbf{\Lambda}} \\ \sqrt{\mathbf{\Lambda}} & \sqrt{1+\mathbf{\Lambda}} \end{pmatrix} \begin{pmatrix} u^{(3)} & \mathbf{0} \\ \mathbf{0} & u^{(4)} \end{pmatrix} \quad (12)$$

where the $u^{(i)}$, ($i = 1, 2, 3, 4$) are $N \times N$ quaternion unitary matrices and $\mathbf{\Lambda}$ is a $N \times N$ real, diagonal, quaternion matrix with non negative elements $\lambda_1, \lambda_2, \dots, \lambda_N$. This parametrization is nothing else than the explicit construction of the symmetric space $\frac{U(N, N, \mathbf{Q})}{U(N, \mathbf{Q}) \otimes U(N, \mathbf{Q})}$, which (through the identification of the quaternion unitary groups as unitary symplectic groups on the complex) is exactly equivalent to the one listed in the fourth line of tab.1. Let us now generalize the DMPK equation to take into account this new critical index. The (generalized) DMPK equation can be obtained as a particular case of the general Calogero-Sutherland construction (see [12,13]). The result is:

$$\frac{\partial P}{\partial s} = \frac{2}{\tilde{\gamma}} \sum_{i=1}^N \frac{\partial}{\partial \lambda_i} \lambda_i (1 + \lambda_i) \tilde{J} \frac{\partial}{\partial \lambda_i} \tilde{J}^{-1} P \quad (14)$$

where

$$\tilde{J}(\{\lambda_n\}) = \prod_{i < j} |\lambda_j - \lambda_i|^\beta \prod_i [\lambda_i (1 + \lambda_i)]^{\frac{\eta-1}{2}} \quad , \quad (15)$$

$\tilde{\gamma} = \frac{2}{\eta+1}(\beta(N-1) + 1 + \eta)$ and the indices β and η are listed in tab.1.

The DMPK equation greatly simplifies if one chooses a new set of variables $\{x_n\}$ related to the eigenvalues $\{\lambda_i\}$ by $\lambda_n = \sinh^2 x_n$. Then the solution can be written in terms of so called ‘‘zonal spherical functions’’ $\Phi_k(x)$,

$x = \{x_1, \dots, x_N\}$, $k = \{k_1, \dots, k_N\}$ (which depend on the chosen symmetric space).

$$P(\{x_n\}, s) = [\xi(x)]^2 \int dk e^{-\frac{k^2}{2\gamma}s} \frac{\Phi_k(x)}{|\Delta(k)|^2} F(k) \quad (16)$$

where

$$\xi(x) = \prod_{i < j} |\sinh^2 x_j - \sinh^2 x_i|^{\frac{\beta}{2}} \prod_i |\sinh 2x_i|^{\frac{\eta}{2}} \quad ; \quad (17)$$

$$F(k) = \prod_j k_j \tanh\left(\frac{\pi k_j}{2}\right) (k_j^2 + 1)^{\frac{\eta-1}{2}} \quad (18)$$

$$|\Delta(k)|^2 = \prod_{m < j} \left| \frac{\Gamma\left(i\frac{k_m - k_j}{2}\right) \Gamma\left(i\frac{k_m + k_j}{2}\right)}{\Gamma\left(\frac{\beta}{2} + i\frac{k_m - k_j}{2}\right) \Gamma\left(\frac{\beta}{2} + i\frac{k_m + k_j}{2}\right)} \right|^2 \quad (19)$$

and Γ denotes the Euler gamma function.

In the insulating regime ($k \ll 1$) we can use the asymptotic expansion described in [12] for the zonal spherical function. In this way the integration over k can be done explicitly and gives [12]:

$$P(\{x_n\}, s) = \prod_{i < j} |\sinh^2 x_j - \sinh^2 x_i|^{\frac{\beta}{2}} [(x_j^2 - x_i^2)] \times \prod_i \left[\exp(-x_i^2 \tilde{\gamma}/(2s)) x_i (\sinh 2x_i)^{\eta/2} \right]. \quad (20)$$

Ordering the x_n 's from small to large and using the fact that in this regime $1 \ll x_1 \ll x_2 \ll \dots \ll x_N$ we can approximate the eigenvalue distribution as follows:

$$P(\{x_n\}, s) = \prod_{i=1}^N \exp[-(\tilde{\gamma}/(2s))(x_i - \bar{x}_i)^2]. \quad (21)$$

where $\bar{x}_n = \frac{s}{\tilde{\gamma}}(\eta + \beta(n-1))$. In the strongly localized limit the conductance is dominated by the first eigenvalue of the transmission matrix:

$$< \log g > \equiv -\frac{2L}{\xi} \sim \log T_1 \sim -2\bar{x}_1 = \frac{2s\eta}{\tilde{\gamma}}. \quad (22)$$

and hence:

$$\xi = \frac{2l}{\eta(\eta+1)} [\beta(N-1) + 1 + \eta] \quad (23)$$

Setting now the proper values of β , N and η for the various regimes, and taking the large N limit, we see, as anticipated, that if both time reversal and Kramers degeneracy are broken, namely if we move from $\frac{SO^*(4N)}{U(2N)} \rightarrow \frac{SU(2N, 2N)}{SU(2N) \otimes SU(2N) \otimes U(1)}$ then ξ does not change, hence $\xi(H)/\xi = 1$ and the MC is 0. If we assume that Kramers degeneracy is conserved then we move from

$\frac{SO^*(4N)}{U(2N)} \rightarrow \frac{USp(2N, 2N)}{USp(2N) \otimes USp(2N)}$ and $\xi(H)/\xi = 1/6$ thus implying a negative MC.

Needless to say that the negative MC measured in the experiments could also simply indicate a failure of the RMT approach outside the metallic phase. Moreover one must be aware of the fact that all the above results are constrained to the quasi one-dimensional limit which underlies the DMPK equation. Nevertheless, it would be very interesting to test the possible existence of this new universality class. In this respect a stringent and simple test is given by the ratio between the mean and the variance of $\log g$. This ratio is easily extracted from eq.s(21,22) and turns out to be

$$\frac{\text{var}(\log g)}{< \log g >} = \frac{2}{\eta}. \quad (24)$$

This observable gives a direct measure of η and could play in the insulating regime the role which is played in the metallic one by the universal conductance fluctuations for β . We hope that our results will stimulate new experiments in order clarify this issue.

-
- [1] E. Medina and M. Kardar, Phys. Rev. Lett. **66**, 3187 (1991); Phys. Rev. B **46**, 9984 (1992).
 - [2] Y. Imry and I. Lerner, preprint cond-mat/9411101
 - [3] J.L. Pichard, M. Sanquer, K. Slevin and P. Debray, Phys. Rev. Lett. **65**, 1812 (1990).
 - [4] Y. Shapir and Z. Ovadyahu, Phys. Rev. **B40**, 12441 (1989); P. Hernandez and M. Sanquer, Phys. Rev. Lett. **68**, 1402 (1992).
 - [5] Y. Meir, N.S. Wingreen, O. Entin-Wohlman and B. Altshuler, Phys. Rev. Lett. **66**, 1517 (1991); Y. Meir and O. Entin-Wohlman; Phys. Rev. Lett. **70**, 1988 (1993);
 - [6] Y. Imry, Europhys. Lett. **1**, 249 (1986).
 - [7] K. A. Muttalib, J.-L. Pichard, and A. D. Stone, Phys. Rev. Lett. **59**, 2475 (1987).
 - [8] K.B. Efetov and A.I. Larkin, Sov. Phys. JETP **58**, 444 (1983); K.B. Efetov, Adv. Phys. **32**, 53 (1983).
 - [9] O. N. Dorokhov, Pis'ma Zh. Eksp. Teor. Fiz. **36**, 259 (1982) [JETP Lett. **36**, 318 (1982)]. P. A. Mello, P. Pereyra, and N. Kumar, Ann. Phys. **181**, 290 (1988). P. A. Mello and A. D. Stone, Phys. Rev. B **44**, 3559 (1991). J. T. Chalker and A. M. S. Macêdo, Phys. Rev. B **46**, 14985 (1992).
 - [10] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces* (Academic, 1978)
 - [11] K. Frahm and J.L. Pichard, preprint cond-mat/9504097
 - [12] M. Caselle, Phys. Rev. Lett. **74** 2776 (1995).
 - [13] M. A. Olshanetsky and A. M. Perelomov, Phys. Rep. **94**, 313 (1984).